

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number
WO 03/104379 A1

(51) International Patent Classification⁷: C11D 3/39, 3/16

(21) International Application Number: PCT/EP03/05217

(22) International Filing Date: 19 May 2003 (19.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0212991.4 6 June 2002 (06.06.2002) GB

(71) Applicant (for AL, AM, AT, AZ, BA, BE, BF, BG, BJ, BR, BY, CF, CG, CH, CI, CM, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GA, GE, GN, GQ, GR, GW, HR, HU, ID, IS, IT, JP, KG, KP, KR, KZ, LR, LT, LU, LV, MA, MC, MD, MG, MK, ML, MR, MX, MZ, NE, NI, NL, NO, PH, PL, PT, RO, RU, SE, SI, SK, SN, TD, TG, TJ, TM, TN, TR, UA, UZ, VN, YU only): UNILEVER N.V. [NL/NL]; UNILEVER N.V., Weena 455, NL-3013 AL Rotterdam (NL).

(71) Applicant (for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, OM, SC, SD, SG, SL, SZ, TT, TZ, UG, VC, ZA, ZM, ZW only): UNILEVER PLC [GB/GB]; UNILEVER HOUSE, Blackfriars, London, Greater London EC4 4BQ (GB).

(71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, 400 020 Mumbai (IN).

(72) Inventors: HAGE, Ronald; UNILEVER R & D VLAARDINGEN, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). LIENKE, Joachim; UNILEVER R & D VLAARDINGEN, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). PETERSEN, Patricia; UNILEVER R & D VLAARDINGEN, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).

(74) Agent: ELLIOTT, Peter, William; UNILEVER PLC, Patent Department, Colworth House, Sharnbrook, Bedfordshire MK44 1LQ (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LIGAND AND COMPLEX FOR CATALYTICALLY BLEACHING A SUBSTRATE

(57) Abstract: The present invention provides a bleaching composition comprising a [3.3.1] bicyclo compound carrying at least one tertiary amine group the bleaching composition substantially devoid of a peroxygen source.

WO 03/104379 A1

LIGAND AND COMPLEX FOR CATALYTICALLY BLEACHING A SUBSTRATEFIELD OF INVENTION

This invention relates to a class of ligand or complex
5 thereof useful as catalysts for catalytically bleaching
substrates in an environment substantially devoid of peroxy
species.

BACKGROUND OF INVENTION

10 The use of bleaching catalysts for stain removal has been
developed over recent years. The recent discovery that some
catalysts are capable of bleaching effectively in the
absence of an added peroxy source has recently become the
focus of some interest, for example: WO9965905; WO0012667;
15 WO0012808; and, WO0029537.


The search for new classes of compounds that are suitable as
"air bleaching" catalysts is ongoing.

20 Various [3.3.1] bicyclo compounds and complexes thereof are
discussed in the literature, see for example: Comba P. et
al., J. Chem. Soc. Dalton Trans, 1998, (23) 3997-4001;
Börzel et al. Chem. Eur. J. 1999, 5, No. 6, 1716 to 1721 and
review by P. Comba in Coordination Chemistry Reviews 2000,
25 200-202, 217 to 245, entitled "Coordination compounds in the
Entactic State". These compounds are discussed in terms of
their physical properties.

WO0060045 discloses a bleaching system comprising: a) from
30 about 1ppb, by weight of a transition metal catalyst

- 2 -

comprising: i) a transition metal; ii) a ligand having formula (I):



wherein each R is independently hydrogen, hydroxyl, C1-C4 alkyl, and mixtures thereof; R1 is C1-C4 alkyl, C6-C10 aryl, and mixtures thereof; R2 is C1-C4 alkyl, C6-C10 aryl, and mixtures thereof; R3 and R4 are each independently hydrogen, C1-C8 alkyl, C1-C8 hydroxyalkyl, $-(CH_2)_xCO_2R_5$ wherein R5 is C1-C4 alkyl, x is from 0 to 4, and mixtures thereof; X is carbonyl, $-C(R_6)_2-$ wherein each R6 is independently hydrogen, hydroxyl, C1-C4 alkyl, and mixtures thereof; b) optionally a source of hydrogen peroxide; and c) the balance carriers and adjunct ingredients. However, the teaching of WO0060045 limits substituents at the nitrogens (3 and 7 positions) of bicyclostructure to homoaromatic carbon groups, namely alkyl and aryl. The general structure of Formula (I) is referred to as a bispidon.

SUMMARY OF INVENTION

The bleaching of a stain by a peroxy species is aided by the presence of an active transition metal catalyst. A peroxy species commonly found in laundry bleaching compositions is hydrogen peroxide (H_2O_2) or a precursor thereof, e.g., sodium percarbonate or sodium perborate. In

- 3 -

many instances an activator/precursor, e.g., TAED (tetraacetylene diamine), is present which serves together with hydrogen peroxide to form a peracid $[RC(O)OOH]$ to facilitate bleaching.

5

Recently we have found that oily stains are bleached in the presence of selected transition metal catalysts in the absence of an added peroxy source. The bleaching of an oily stain in the absence of an added peroxy source has been attributed to oxygen derived from the air. Whilst it is true that bleaching is effected by oxygen sourced from the air the route in which oxygen plays a part is becoming understood. In this regard, the term "air bleaching" is used.

15

We have concluded from our research that bleaching of a chromophore in an oily stain is effected by products formed by adventitious oxidation of components in the oily stain. These products, alkyl hydroperoxides, are generated naturally by autoxidation of the oily stain and the alkyl hydroperoxides together with a transition metal catalyst serve to bleach chromophores in the oily stain. Alkyl hydroperoxides ($ROOH$) are generally less reactive than other peroxy species, for example, peracids $(RC(O)OOH)$, hydrogen peroxide (H_2O_2), percarbonates and perborates.

20

Our earlier filed application PCT/EP01/13314, filed 15 November 2002, which claims priority from GB0030673.8, filed 15 December 2000, discloses the use of various bispidon compounds. Referring to the structure above, PCT/EP01/13314 teaches that there is an advantage to be secured by having

30


- 4 -

at least one of R1 and R2 as group containing a heteroatom capable of coordinating to a transition metal. We have now found that by having at least one of R1 and R2 as a group that is a tertiary amine linked to one or more of the nitrogen atoms of the bicyclo structure by a C2 to C4 alkyl chain further advantages are secured. In addition, we have also found that heterocycles other than pyridyl may be used at the 2 and 4 positions.

- 10 Accordingly, in a first aspect, the present invention provides a bleaching composition comprising:

A bleaching composition comprising:

- 15 a) a monomer ligand, L, or transition metal catalyst thereof of a ligand having the formula (I):



- 20 wherein at least one of R1 and R2 is an optionally substituted tertiary amine of the form -C2-C4-alkyl-NR7R8, in which R7 and R8 are independently selected from the group consisting of straight chain, branched or cyclo C1-C12 alkyl, benzyl, the -C2-C4-alkyl- of the -C2-C4-alkyl-NR7R8 may be substituted by 1 to 4 C1-C2-alkyl, or may form part of a C3 to C6 alkyl ring, and in which R7 and R8 may
- 25

- 5 -

together form a saturated ring containing one or more other heteroatoms, the other of R1 and R2 being independently selected from:

-C2-C4-alkyl-NR7R8 as defined above,

5 -C1-C24-optionally substituted-alkyl,

-C6-C10-aryl, -C1-C4-alkyl-C6-C10-aryl,

a heterocycloalkyl: selected from the group consisting of:

pyrrolinyl, pyrrolidinyl, morpholinyl, piperidinyl,

piperazinyl, hexamethylene imine, 1,4-piperazinyl,

10 tetrahydrothiophenyl, tetrahydrofuran, tetrahydropyran, and oxazolidinyl, wherein the heterocycloalkyl may be

connected to the ligand via any atom in the ring of the selected heterocycloalkyl,

a -C1-C6-alkyl-heterocycloalkyl, wherein the

15 heterocycloalkyl of the -C1-C6-heterocycloalkyl is selected

from the group consisting of: piperidinyl, piperidine, 1,4-piperazine, tetrahydrothiophene, tetrahydrofuran,

pyrrolidine, and tetrahydropyran, wherein the

heterocycloalkyl may be connected to the -C1-C6-alkyl via

20 any atom in the ring of the selected heterocycloalkyl,

a -C1-C6-alkyl-heteroaryl, wherein the heteroaryl of the -

C1-C6-alkylheteroaryl is selected from the group consisting of: pyridinyl, pyrimidinyl, pyrazinyl, triazolyl,

pyridazinyl, 1,3,5-triazinyl, quinolinyl, isoquinolinyl,

25 quinoxalinyl, imidazolyl, pyrazolyl, benzimidazolyl,

thiazolyl, oxazolidinyl, pyrrolyl, carbazolyl, indolyl, and

isoindolyl, wherein the heteroaryl may be connected to the -

C1-C6-alkyl via any atom in the ring of the selected

heteroaryl and the selected heteroaryl is optionally

30 substituted by -C1-C4-alkyl, -C0-C6-alkyl-phenol, -C0-C6-

- 6 -

alkyl-thiophenol, -C2-C4-alkyl-thiol, -C2-C4-alkyl-thioether, -C2-C4-alkyl-alcohol, -C2-C4-alkyl-amine, and a -C2-C4-alkyl-carboxylate;

- 5 R3 and R4 are independently selected from hydrogen, C1-C4-alkyl, phenyl, electron withdrawing groups and reduced products and derivatives thereof;

10 X is selected from: C=O, a ketal derivative of C=O, a thioketal of derivative of C=O, and $-[C(R6)_2]_y-$ wherein y takes a value 0 or 1; each R6 is independently selected from hydrogen, hydroxyl, O-C1-C24-alkyl, O-benzyl, O-(C=O)-C1-C24-alkyl, C1-C24-alkyl;

15 z groups are same heteroaromatic groups, selected from the group consisting of: pyridinyl; pyrimidinyl; pyrazinyl; triazolyl; pyridazinyl; 1,3,5-triazinyl; quinolinyl; isoquinolinyl; quinoxalinyl; imidazolyl; pyrazolyl; benzimidazolyl; thiazolyl; oxazolidinyl; pyrrolyl; 20 carbazolyl; indolyl; and isoindolyl, and the selected Z is optionally substituted by -C1-C4-alkyl;

b) the balance carriers and adjunct ingredients.

25 In a second aspect, the present invention provides a bleaching composition comprising, in an aqueous medium, the bicyclo ligand of the general Formula (I) which forms a complex with a transition metal, the complex catalysing bleaching of a substrate, wherein the aqueous medium is 30 substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system. It is preferred that

- 7 -

the medium has a pH value in the range from pH 6 to 12 and most preferably from pH 8 to 11.

5 The term "substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system" should be construed within spirit of the invention. It is preferred that the composition has as low a content of peroxy species present as possible. It is preferred that the bleaching formulation contains less than 1 % wt/wt total concentration
10 of peracid or hydrogen peroxide or source thereof, preferably the bleaching formulation contains less than 0.3 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, most preferably the bleaching composition is devoid of peracid or hydrogen peroxide or source thereof.

15 An advantage of the class of ligand and complex according to the present invention is that the complex can catalyse bleaching of a substrate via atmospheric oxygen, thus permitting its use in a medium such as an aqueous medium
20 that is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system. We have also found that complexes of this class are surprisingly effective in catalysing bleaching of the substrate via atmospheric oxygen after treatment of the substrate.

25 One skilled in the art will appreciate that not all peroxy activating catalysts are capable of providing discernable bleaching activity with adventitious hydroperoxides present in a stain. However, the converse is not true. There is no
30 evidence to indicate that any "air bleaching" catalyst will not function as peroxy activating catalyst. In this

- 8 -

regard, all "air bleaching" catalysts disclosed herein may be used as a peroxy activating catalyst. Catalysts of the present invention may be incorporated into a composition together with a peroxy species or source thereof. For a
5 discussion of acceptable ranges of a peroxy species or source thereof and other adjuvants that may be present the reader is directed to United States Patent 6,022,490, the contents of which are incorporated by reference.

10 The present invention extends to a method of bleaching a substrate comprising applying to the substrate, in an aqueous medium, the bleaching composition according to the present invention.

15 The present invention extends to a commercial package comprising the bleaching composition according to the present invention together with instructions for its use.

The present invention further provides a dry textile having
20 an organic substance as defined above applied or deposited thereon, whereby bleaching via atmospheric oxygen derived alkylhydroperoxides is catalysed on the textile.

Advantageously, by enabling a bleaching effect even after
25 the textile has been treated, the benefits of bleaching can be prolonged on the textile. Furthermore, since a bleaching effect is conferred to the textile after the treatment, the treatment itself, such as a laundry wash cycle, may for example be shortened. Moreover, since a bleaching effect is
30 achieved via atmospheric oxygen after treatment of the

- 9 -

textile, hydrogen peroxide or peroxy-based bleach systems can be omitted from the treatment substance.

5 The organic substance may be contacted to the textile fabric in any suitable manner. For example, it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example as an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid.

10 Other suitable means of contacting the organic substance to the textile may be used, as further explained below.

Any suitable textile that is susceptible to bleaching or one that one might wish to subject to bleaching may be used.

15 Preferably the textile is a laundry fabric or garment.

In a preferred embodiment, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures. The elevated temperatures are commonly

20 provided by a heated agitated environment, as for example found in a tumble dryer, which has been found to accelerate and enhance the "air bleaching" effect.

The organic substance can be contacted with the textile

25 fabric in any conventional manner. For example it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example in an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on

30 aerosol fluid.

- 10 -

In a particularly preferred embodiment the method according to the present invention is carried out on a laundry fabric using aqueous treatment liquor. In particular the treatment may be effected in, or as an adjunct to, an essentially
5 conventional wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent wash liquor. The organic substance can be delivered into the wash liquor from a powder, granule, pellet, tablet, block, bar or other such solid form. The
10 solid form can comprise a carrier, which can be particulate, sheet-like or comprise a three-dimensional object. The carrier can be dispersible or soluble in the wash liquor or may remain substantially intact. In other embodiments, the organic substance can be delivered into the wash liquor from
15 a paste, gel or liquid concentrate.

It is particularly advantageous that the organic substance used in the method of the present invention makes use of atmospheric oxygen in its bleaching activity. This avoids
20 the requirement that peroxygen bleaches and/or other relatively large quantities of reactive substances need be used in the treatment process. Consequently, only a relatively small quantity of bleach active substance need be employed and this allows dosage routes to be exploited that
25 could previously not be used. Thus, while it is preferable to include the organic substance in a composition that is normally used in a washing process, such as a pre-treatment, main-wash, conditioning composition or ironing aid, other means for ensuring that the organic substance is present in
30 the wash liquor may be envisaged.

- 11 -

For example, it is envisaged that the organic substance can be presented in the form of a body from which it is slowly released during the whole or part of the laundry process. Such release can occur over the course of a single wash or
5 over the course of a plurality of washes. In the latter case it is envisaged that the organic substance can be released from a carrier substrate used in association with the wash process, e.g. from a body placed in the dispenser drawer of
10 a washing machine, elsewhere in the delivery system or in the drum of the washing machine. When used in the drum of the washing machine the carrier can be freely moving or fixed relative to the drum. Such fixing can be achieved by mechanical means, for example by barbs that interact with the drum wall, or employ other forces, for example a
15 magnetic force. The modification of a washing machine to provide for means to hold and retain such a carrier is envisaged similar means being known from the analogous art of toilet block manufacture. Freely moving carriers such as shuttles for dosage of surfactant materials and/or other
20 detergent ingredients into the wash can comprise means for the release of the organic substance into the wash.

In the alternative, the organic substance can be presented in the form of a wash additive that preferably is soluble.
25 The additive can take any of the physical forms used for wash additives, including powder, granule, pellet, sheet, tablet, block, bar or other such solid form or take the form of a paste, gel or liquid. Dosage of the additive can be unitary or in a quantity determined by the user. While it is
30 envisaged that such additives can be used in the main

- 12 -

washing cycle, the use of them in the conditioning or drying cycle is not hereby excluded.

The present invention is not limited to those circumstances
5 in which a washing machine is employed, but can be applied where washing is performed in some alternative vessel. In these circumstances it is envisaged that the organic substance can be delivered by means of slow release from the bowl, bucket or other vessel which is being employed, or
10 from any implement which is being employed, such as a brush, bat or dolly, or from any suitable applicator.

Suitable pre-treatment means for application of the organic substance to the textile material prior to the main wash
15 include sprays, pens, roller-ball devices, bars, soft solid applicator sticks and impregnated cloths or cloths containing microcapsules. Such means are well known in the analogous art of deodorant application and/or in spot treatment of textiles. Similar means for application are
20 employed in those embodiments where the organic substance is applied after the main washing and/or conditioning steps have been performed, e.g. prior to or after ironing or drying of the cloth. For example, the organic substance may be applied using tapes, sheets or sticking plasters coated
25 or impregnated with the substance, or containing microcapsules of the substance. The organic substance may for example be incorporated into a drier sheet so as to be activated or released during a tumble-drier cycle, or the substance can be provided in an impregnated or microcapsule-
30 containing sheet so as to be delivered to the textile when ironed.

- 13 -

Many transition metal complexes have high extinction coefficients in the visible. In this regard, use over time may result in some colour deposition on a substrate after repeated washing. The addition of a limited amount of a
5 peroxy source serves to reduce colour deposition in those instances in which it occurs whilst still permitting "air bleaching". Nevertheless, we have found that in certain instances the free ligand may be used in the bleaching composition of the present invention. By using a free
10 ligand, a bleaching formulation may be prepared that is consistent with consumer formulation colour expectation. In such a formulation the metal ion may be provided by the composition or by trace metals found in the stain.


15 DETAILED DESCRIPTION OF THE INVENTION

The ligand as described herein is capable of dynamic inversion. The ability of the ligand to chelate to a TM depends upon the stereochemistry of the substituents. It is preferred that substituents are endo-endo, but it is likely
20 that stereochemical conversion takes place by retro-Mannich conversion. Retro-Mannich may be prevented by changing the groups present such that retro-Mannich reactions are unfavoured. Nevertheless, it is likely that endo-exo and exo-exo ligands as described herein coordinate to transition
25 metal ions in many instances and are capable of functioning as air bleaching catalysts.

Referring to ligands and complexes thereof and bleaching compositions derived therefrom with respect to Formula (I),
30 it is preferred that each z is the same; and $R_3 = R_4$. Preferred Z groups are pyridine, benzimidazole, thiazole,

- 14 -

imidazole. Most preferred z groups are of the form



, wherein R is independently selected from:


hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkyl-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl. Of the R groups it is preferred that R is H or -C1-C4-alkyl.


It is preferred that R₃ and R₄ are selected from the group consisting of: -CH₂OH, -CH₂OC(O)C1-C₂₀-alkyl, -C(O)O-C1-C₆-alkyl, benzyl ester, -CN, C1-C₆-alkyl, benzyl, phenyl, and C1-C₄-OR wherein R is selected from the group consisting of H, C1-C₂₄-alkyl or C(O)-C1-C₂₄-alkyl. Most preferred R₃ and R₄ groups are: -C(O)-O-CH₃, -C(O)-O-CH₂CH₃, benzyl ester, and CH₂OH. Notwithstanding the above, R₃ and R₄ are initially important such that synthesis proceeds to provide the core structure of Formula (I), namely the [3.3.1] bicyclo structure. After the core structure of Formula (I) is formed reduction or other synthetic methods may change R₃ and R₄. Whilst R₃ and R₄ may participate in influencing the activity of the catalyst in the broadest aspect R₃ and R₄ may be independently selected from electron withdrawing groups and reduced products and derivatives thereof. The R₃ and R₄ substituents do not have a substantial influence on the catalyst per se except to change the hydrophobisity/solubility of the ligand or transition metal catalyst formed therefrom.

It is preferred that the -C₂-C₄-alkyl-NR₇R₈ is selected from the group consisting of: -CH₂CH₂-NR₇R₈, -CH₂CMe₂-NR₇R₈, -CMe₂CH₂-NR₇R₈, -CMeHCH₂-NR₇R₈, -CMeH-CMeH-NR₇R₈, -CH₂CMeH-

- 15 -








NR7R8, -CH₂CH₂CH₂-NR7R8, -CH₂CH₂CMe₂-NR7R8, -CH₂CMe₂CH₂-

NR7R8, -CH₂CH₂-NEt₂, -CH₂CH₂-N(*i*-Pr)₂, , and



, . Preferred groups for R7 and R8 are CH₃, -C₂H₅, -C₃H₇, -C₄H₉, -C₅H₁₁, -C₆H₁₃, and -CH₂C₆H₅. It is most preferred that at least one of R7 and R8 is an optionally substituted alkyl chain of at least five carbon atoms.

Other preferred groups for R7 and R8 are -CH₃, -CH₂CH₃, -CH(CH₃)₂ or where R7 and R8 together with the N form a optionally substituted cyclic structure selected from the

group consisting of: , , , , , , and . Of the


R1 and R2 it is most preferred that R1 is a C₂-C₄-alkyl-NR7R8 and most preferably both R1 and R2 are independently C₂-C₄-alkyl-NR7R8. It is preferred that the -C₂-C₄-alkyl-NR7R8 is a C₂-alkyl-NR7R8.

It is preferred that X is selected from: C=O, and -[C(R₆)₂] wherein each R₆ is independently selected from hydrogen, hydroxyl, C₁-C₂₄-alkoxy and C₁-C₂₄-alkyl. In particular X is preferred in the form C(OH)₂, *syn*-CH(OH) and *anti*-CH(OH). One


- 16 -

skilled in the art will be aware that when $X = C(OH)_2$ a solvent adduct may exist, e.g., when present in methanol the $C(OMe)_2$ adduct will be formed. This adduct will exchange in water such that $X = C(OH)_2$ which again is equilibrium with $X = C=O$.

Particularly preferred ligands are exemplified below:



wherein $-NR_6R_7$ is selected from the group consisting of -




The catalyst may be used as a preformed complex of the ligand and a transition metal. Alternatively, the catalyst may be formed from the free ligand that complexes with a transition metal already present in the water or that complexes with a transition metal present in the substrate. The composition may also be formulated as a composition of the free ligand or a transition metal-substitutable metal-ligand complex, and a source of transition metal, whereby the complex is formed *in situ* in the medium.

- 17 -

The ligand forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI, vanadium II-V and molybdenum II-VI.

The ligand forms a complex of the general formula (A1):



in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI), preferably selected from Fe(II)-(III)-(IV)-(V);

L represents a ligand as herein defined, or its protonated or deprotonated analogue;

X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner, preferably selected from O^{2-} , RBO_2^{2-} , $RCOO^-$, $RCONR^-$, OH^- , NO_3^- , NO , S^{2-} , RS^- , PO_4^{3-} , PO_3OR^{3-} , H_2O , CO_3^{2-} , HCO_3^- , ROH , $N(R)_3$, ROO^- , O_2^{2-} , O_2^- , RCN , Cl^- , Br^- , OCN^- , SCN^- , CN^- , N_3^- , F^- , I^- , RO^- , ClO_4^- , and $CF_3SO_3^-$, and more preferably selected from O^{2-} , RBO_2^{2-} , $RCOO^-$, OH^- , NO_3^- , S^{2-} , RS^- , PO_3^{4-} , H_2O , CO_3^{2-} , HCO_3^- , ROH , $N(R)_3$, Cl^- , Br^- , OCN^- , SCN^- , RCN , N_3^- , F^- , I^- , RO^- , ClO_4^- , and $CF_3SO_3^-$;

Y represents any non-coordinated counter ion, preferably selected from ClO_4^- , BR_4^- , $[MX_4]^-$, $[MX_4]^{2-}$, PF_6^- ,

- 18 -

RCOO^- , NO_3^- , RO^- , $\text{N}^+(\text{R})_4$, ROO^- , O_2^{2-} , O_2^- , Cl^- , Br^- , F^- , I^- , CF_3SO_3^- , $\text{S}_2\text{O}_6^{2-}$, OCN^- , SCN^- , H_2O , RBO_2^{2-} , BF_4^- and BPh_4^- , and more preferably selected from ClO_4^- , BR_4^- , $[\text{FeCl}_4]^-$, PF_6^- , RCOO^- , NO_3^- , RO^- , $\text{N}^+(\text{R})_4$, Cl^- , Br^- , F^- , I^- , CF_3SO_3^- , $\text{S}_2\text{O}_6^{2-}$,

5 OCN^- , SCN^- , H_2O and BF_4^- ;

a represents an integer from 1 to 10, preferably from 1 to 4;

k represents an integer from 1 to 10;

n represents an integer from 1 to 10, preferably from 1 to 4;

m represents zero or an integer from 1 to 20, preferably from 1 to 8; and

each R independently represents a group selected from hydrogen, hydroxyl, $-\text{R}'$ and $-\text{OR}'$, wherein $\text{R}' = \text{alkyl}$, alkenyl ,
 15 cycloalkyl , heterocycloalkyl , aryl , heteroaryl or a carbonyl derivative group, R' being optionally substituted by one or more functional groups E, wherein E independently represents a functional group selected from $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{OH}$, $-\text{OR}'$, $-\text{NH}_2$, $-\text{NHR}'$, $-\text{N}(\text{R}')_2$, $-\text{N}(\text{R}')_3^+$, $-\text{C}(\text{O})\text{R}'$, $-\text{OC}(\text{O})\text{R}'$, $-\text{COOH}$, $-\text{COO}^-$
 20 $(\text{Na}^+, \text{K}^+)$, $-\text{COOR}'$, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NHR}'$, $-\text{C}(\text{O})\text{N}(\text{R}')_2$, heteroaryl , $-\text{R}'$, $-\text{SR}'$, $-\text{SH}$, $-\text{P}(\text{R}')_2$, $-\text{P}(\text{O})(\text{R}')_2$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{P}(\text{O})(\text{OR}')_2$, $-\text{NO}_2$, $-\text{SO}_3\text{H}$, $-\text{SO}_3^-(\text{Na}^+, \text{K}^+)$, $-\text{S}(\text{O})_2\text{R}'$, $-\text{NHC}(\text{O})\text{R}'$, and $-\text{N}(\text{R}')\text{C}(\text{O})\text{R}'$, wherein R' represents cycloalkyl , aryl , arylalkyl , or alkyl optionally substituted by $-\text{F}$, $-\text{Cl}$, $-\text{Br}$,
 25 $-\text{I}$, $-\text{NH}_3^+$, $-\text{SO}_3\text{H}$, $-\text{SO}_3^-(\text{Na}^+, \text{K}^+)$, $-\text{COOH}$, $-\text{COO}^-(\text{Na}^+, \text{K}^+)$, $-\text{P}(\text{O})(\text{OH})_2$, or $-\text{P}(\text{O})(\text{O}^-(\text{Na}^+, \text{K}^+))_2$, and preferably each R independently represents hydrogen, optionally substituted alkyl or optionally substituted aryl , more preferably hydrogen or optionally substituted phenyl , naphthyl or C_{1-4} -
 30 alkyl .

- 19 -

The counter ions Y in formula (A1) balance the charge z on the complex formed by the ligand L, metal M and coordinating species X. Thus, if the charge z is positive, Y may be an anion such as RCOO^- , BPh_4^- , ClO_4^- , BF_4^- , PF_6^- , RSO_3^- , RSO_4^- , SO_4^{2-} , NO_3^- , F^- , Cl^- , Br^- , or I^- , with R being hydrogen, optionally substituted alkyl or optionally substituted aryl. If z is negative, Y may be a common cation such as an alkali metal, alkaline earth metal or (alkyl)ammonium cation.

Suitable counter ions Y include those which give rise to the formation of storage-stable solids. Preferred counter ions for the preferred metal complexes are selected from R^7COO^- , ClO_4^- , BF_4^- , PF_6^- , RSO_3^- (in particular CF_3SO_3^-), RSO_4^- , SO_4^{2-} , NO_3^- , F^- , Cl^- , Br^- , and I^- , wherein R represents hydrogen or optionally substituted phenyl, naphthyl or $\text{C}_1\text{-C}_4$ alkyl.

The novel compounds of Formula (I) as provided by the present invention also extend to their various transition metal complexes, the transition metal complexes are as discussed above with reference to (A1).

It will be appreciated that the complex (A1) can be formed by any appropriate means, including *in situ* formation whereby precursors of the complex are transformed into the active complex of general formula (A1) under conditions of storage or use. Preferably, the complex is formed as a well-defined complex or in a solvent mixture comprising a salt of the metal M and the ligand L or ligand L-generating species. Alternatively, the catalyst may be formed *in situ* from suitable precursors for the complex, for example in a solution or dispersion containing the precursor materials.

- 20 -

In one such example, the active catalyst may be formed *in situ* in a mixture comprising a salt of the metal M and the ligand L, or a ligand L-generating species, in a suitable solvent. Thus, for example, if M is iron, an iron salt such as FeSO_4 can be mixed in solution with the ligand L, or a ligand L-generating species, to form the active complex. Thus, for example, the composition may be formed from a mixture of the ligand L and a metal salt MX_n in which preferably $n=1-5$, more preferably 1-3. In another such example, the ligand L, or a ligand L-generating species, can be mixed with metal M ions present in the substrate or wash liquor to form the active catalyst *in situ*. Suitable ligand L-generating species include metal-free compounds or metal coordination complexes that comprise the ligand L and can be substituted by metal M ions to form the active complex according to the formula (A1).

The catalysts according to the present invention may be used for laundry cleaning, hard surface cleaning (including cleaning of lavatories, kitchen work surfaces, floors, mechanical ware washing etc.). As is generally known in the art, bleaching compositions are also employed in waste-water treatment, pulp bleaching during the manufacture of paper, leather manufacture, dye transfer inhibition, food processing, starch bleaching, sterilisation, whitening in oral hygiene preparations and/or contact lens disinfection.

In typical washing compositions the level of the organic substance is such that the in-use level is from 1 μM to 50mM, with preferred in-use levels for domestic laundry operations falling in the range 10 to 100 μM . Higher levels may be

- 21 -

desired and applied in industrial bleaching processes, such as textile and paper pulp bleaching. These levels reflect the amount of catalyst that may be present in a wash dose of a detergent composition. The bleaching composition
5 comprises at least 1 ppb of the ligand or complex thereof.

In the context of the present invention, bleaching should be understood as relating generally to the decolourisation of stains or of other materials attached to or associated with
10 a substrate. However, it is envisaged that the present invention can be applied where a requirement is the removal and/or neutralisation by an oxidative bleaching reaction of malodours or other undesirable components attached to or otherwise associated with a substrate. Furthermore, in the
15 context of the present invention bleaching is to be understood as being restricted to any bleaching mechanism or process that does not require the presence of light or activation by light.


20 Synthesis

In addition to the utility of the ligands and complexes of the present invention as catalysts another advantage is that the ligands are generally relatively easy to synthesize in comparison to other ligands. The following is one example
25 of a strategic synthetic approach; it will be evident to one skilled in the art of synthetic organic chemistry that many approaches may be taken to obtain ligands and complexes for use in the present invention. The ease of synthesis of the ligand of Formula (I) is dependent upon the nature of
30 substituents about the structure. The ligands of Formula (I) are most preferably symmetric. Synthesis of these types of


- 22 -

molecules are found in articles by U. Holzgrabe et al. in Arch. Pharm. (Weinheim, Ger.) 1992, 325, 657 and A. Samhammer et al. Arch. Pharm. (Weinheim, Ger.) 1984, 322, 557. Below is given a schematic example illustrating the ease of synthesis. The synthesis is shown in a two step synthesis, Scheme 1 and Scheme 2, but in some cases may be conducted as a "one-pot" synthesis depending upon the nature of the substituents. Nevertheless, where substituents at positions 7 and 3 are different a two step synthesis is preferred. The product of reaction as found in Scheme 1 is referred to as dimethyl 2,6-di-(2-pyridyl)-1-methyl-piperid-4-one-3,5-dicarboxylate (NPy2), which can easily tautomerize to the enol. The synthesis is exemplified in R. Haller, K.W. Merz, *Pharm. Acta Helv.*, **1963**, 442.

Scheme 1



Scheme 2




Another intermediate that may be produced according to the general teachings of Scheme 1 wherein MeNH₂ is replaced by

- 23 -

Me₂NCH₂CH₂NH₂ such that a product referred to as dimethyl-2,6-di-(2-pyridyl)-1-(N,N-dimethylamino)ethylene-piperid-4-one-3,5-dicarboxylate is produced, the structure of which is given below.

5



One skilled in the art will appreciate that whilst Ac [-CO(O)Me] is an electron withdrawing group and electron withdrawing groups are generally preferred to facilitate synthesis other groups will also allow the reaction to proceed. Examples of suitable electron withdrawing groups are given above and will be evident to one skilled in the art. The reaction is also driven by precipitation of the product from solution.

15

In instances, depending upon the nature of the substituents, for example a phenolic group, it will be necessary to protect certain functional groups. The choice of protecting groups during synthesis to prevent undesirable reactions will be evident to one skilled in the art. For a discussion of protecting groups in organic synthesis the reader is directed to T. W. Green and P. G. M. Wuts, Protective Groups In Organic Synthesis 3rd Ed.; J. Wiley and Sons, 1999.


20

25

It will be evident that if a diamine is substituted for methylamine in the reaction illustrated in Scheme 2 two

- 24 -

structures may be linked together via the 7 positions as found in the structure below.



5 In addition, if a diamine is substituted for methylamine in the reaction illustrated in Scheme 1 a NPy2 structure is formed that is linked at the 3 positions. Obviously, this dimer would serve as a precursor to other dimer and polymer type structures. The present invention is confined to
 10 "monomer" ligands and not the dimer and polymer units linked by a covalent bond as described above. The term "monomer" as used herein is used to exclude these products in which covalently linked polyligand type structures are formed.

15 The Detergent Composition.

The air bleach catalyst and may be used in a detergent composition specifically suited for stain bleaching purposes, and this constitutes a second aspect of the invention. To that extent, the composition comprises a
 20 surfactant and optionally other conventional detergent ingredients. The invention in its second aspect provides an enzymatic detergent composition which comprises from 0.1 - 50 % by weight, based on the total detergent composition, of one or more surfactants. This surfactant system may in turn
 25 comprise 0 - 95 % by weight of one or more anionic

- 25 -

surfactants and 5 to 100 % by weight of one or more nonionic surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost.

- 5 The enzymatic detergent composition according to the invention will generally be used as a dilution in water of about 0.05 to 2%.

The condition of "the balance carriers and adjunct
10 ingredients" should be taken to be at least 1% wt/wt of a surfactant, preferably at least 5% wt/wt.

In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants
15 described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H.
20 Stache, 2nd Edn., Carl Hauser Verlag, 1981.

Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for
25 example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are C₆-C₂₂ alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide
30 per molecule, and the condensation products of aliphatic C₈-

- 26 -

C₁₈ primary or secondary linear or branched alcohols with ethylene oxide, generally 5 to 40 EO.

Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C₈-C₁₈ alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C₉-C₂₀ benzene sulphonates, particularly sodium linear secondary alkyl C₁₀-C₁₅ benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C₁₁-C₁₅ alkyl benzene sulphonates and sodium C₁₂-C₁₈ alkyl sulphates. Also applicable are surfactants such as those described in EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system that is a mixture of an alkali metal salt of a C₁₆-C₁₈ primary alcohol sulphate together with a C₁₂-C₁₅ primary alcohol 3-7 EO ethoxylate.

- 27 -

The nonionic detergent is preferably present in amounts greater than 10%, e.g. 25-90% by weight of the surfactant system. Anionic surfactants can be present for example in amounts in the range from about 5% to about 40% by weight of the surfactant system.

The detergent composition may take any suitable physical form, such as a powder, granular composition, tablets, a paste or an anhydrous gel.

Enzymes

The detergent compositions of the present invention may additionally comprise one or more enzymes, which provide cleaning performance, fabric care and/or sanitation benefits.

Said enzymes include oxidoreductases, transferases, hydrolases, lyases, isomerases and ligases. Suitable members of these enzyme classes are described in Enzyme nomenclature 1992: recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology on the nomenclature and classification of enzymes, 1992, ISBN 0-12-227165-3, Academic Press.

Examples of the hydrolases are carboxylic ester hydrolase, thiolester hydrolase, phosphoric monoester hydrolase, and phosphoric diester hydrolase which act on the ester bond; glycosidase which acts on O-glycosyl compounds; glycosylase hydrolysing N-glycosyl compounds; thioether hydrolase which acts on the ether bond; and exopeptidases and endopeptidases which act on the peptide bond. Preferable among them are

- 28 -

carboxylic ester hydrolase, glycosidase and exo- and endopeptidases. Specific examples of suitable hydrolases include (1) exopeptidases such as aminopeptidase and carboxypeptidase A and B and endopeptidases such as pepsin, 5 pepsin B, chymosin, trypsin, chymotrypsin, elastase, enteropeptidase, cathepsin B, papain, chymopapain, ficain, thrombin, plasmin, renin, subtilisin, aspergillopepsin, collagenase, clostripain, kallikrein, gastricsin, cathepsin D, bromelain, chymotrypsin C, urokinase, cucumisin, oryzin, 10 proteinase K, thermomycolin, thermitase, lactocepin, thermolysin, bacillolysin. Preferred among them is subtilisin; (2) glycosidases such as α -amylase, β -amylase, glucoamylase, isoamylase, cellulase, endo-1,3(4)- β -glucanase (β -glucanase), xylanase, dextranase, polygalacturonase 15 (pectinase), lysozyme, invertase, hyaluronidase, pullulanase, neopullulanase, chitinase, arabinosidase, exocellobiohydrolase, hexosaminidase, mycodextranase, endo-1,4- β -mannanase (hemicellulase), xyloglucanase, endo- β -galactosidase (keratanase), mannanase and other saccharide 20 gum degrading enzymes as described in WO-A-99/09127. Preferred among them are α -amylase and cellulase; (3) carboxylic ester hydrolase including carboxylesterase, lipase, phospholipase, pectinesterase, cholesterol esterase, chlorophyllase, tannase and wax-ester hydrolase. Preferred 25 among them is lipase.

Examples of transferases and ligases are glutathione S-transferase and acid-thiol ligase as described in WO-A-98/59028 and xyloglycan endotransglycosylase as 30 described in WO-A-98/38288.

- 29 -

Examples of lyases are hyaluronate lyase, pectate lyase, lipex, chondroitinase, pectin lyase, alginase II. Especially preferred is pectolyase, which is a mixture of pectinase and pectin lyase.

5

Examples of the oxidoreductases are oxidases such as glucose oxidase, methanol oxidase, bilirubin oxidase, catechol oxidase, laccase, peroxidases such as ligninase and those described in WO-A-97/31090, monooxygenase, dioxygenase such as lipoygenase and other oxygenases as described in WO-A-99/02632, WO-A-99/02638, WO-A-99/02639 and the cytochrome based enzymatic bleaching systems described in WO-A-99/02641.

10

15 The activity of oxidoreductases, in particular the phenol oxidising enzymes in a process for bleaching stains on fabrics and/or dyes in solution and/or antimicrobial treatment can be enhanced by adding certain organic compounds, called enhancers. Examples of enhancers are 2,2'-
20 azo-bis-(3-ethylbenzo-thiazoline-6-sulphonate (ABTS) and Phenothiazine-10-propionate (PTP). More enhancers are described in WO-A-94/12619, WO-A-94/12620, WO-A-94/12621, WO-A-97/11217, WO-A-99/23887. Enhancers are generally added at a level of 0.01% to 5% by weight of detergent
25 composition.

Builders, polymers and other enzymes as optional ingredients may also be present as found in WO0060045.

30

Suitable detergency builders as optional ingredients may also be present as found in WO0034427.

- 30 -

The invention will now be further illustrated by way of the following non-limiting examples:

EXAMPLES

- 5 The ligand N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane (MeN4py) was prepared as described in EP 0 909 809 A2. The synthesis of the iron complex, [(MeN4Py)FeCl]Cl, has been described elsewhere (WO 0116271).

10 **Synthesis**

Procedure for bispidone synthesis:

- A suspension of 7.15 g (16.3 mmol) of piperidone (Npy2) (synthesis exemplified in R. Haller, K.W. Merz, *Pharm. Acta Helv.*, **1963**, 442) in 40 ml ethanol is treated with 1.72 g (19.6 mmol) of N,N-dimethylethylenediamine and 3.5 ml of formaldehyde (37% in water) - 36.1 mmol) and is refluxed for 30 min. The resulting clear, slight yellow to dark brown reaction solution is evaporated to half of its volume and
20 left at 5°C for 24 h. The yellow precipitate formed is filtered, washed with little EtOH until the precipitate is white and dried under high vacuum.

- If no precipitate is obtained, the reaction mixture is
25 evaporated to dryness, dissolved in as little EtOH as possible and left at 5°C for 72 h.

- 31 -

Analytical data:**Melting point:** 147°C.**5 CHN analysis:**

calc. (%) C 63.02 H 6.71 N 14.13

found (%) C 62.69 H 6.76 N 13.79

FAB⁺MS (NBA): 496.3 (MH⁺); C₂₆H₃₃N₅O₅ M = 495.25 g/mol

10

IR[cm⁻¹]: 3039 (w), 2942 (m), 2779 (m), 2760 (m), 2708 (w),
1723 (s), 1587 (m), 1465 (m), 1431(m), 1270 (s), 1162 (m),
971 (m), 751 (m).

15 **¹H-NMR** (300.133 MHz, CDCl₃): δ = 1.98 (s, 3H, N-CH₃), 2.32
(bs, 6H, N-(CH₃)₂), 2.49 (bs, 4H, N-CH₂-), 2.61 (d, 2H, ²J_{HH} =
12.1 Hz, -CH₂-), 3.12 (d, 2H, ²J_{HH} = 9.5 Hz, -CH₂-), 3.79 (s,
6H, OCH₃), 4.66 (s, 2H, CH-Py), 7.20 (dt, 2H, ³J_{HH} = 4.8 Hz,
⁴J_{HH} = 1.1 Hz, Py-H), 7.73 (dt, 2H, ³J_{HH} = 7.7 Hz, ⁴J_{HH} = 1.8
20 Hz, Py-H), 8.11 (bd, 2H, ³J_{HH} = 7.7 Hz, Py-H), 8.47 (dd, 2H,
³J_{HH} = 8.5 Hz, ⁴J_{HH} = 1.1 Hz, Py-H).

¹³C-NMR (75.47 MHz, CDCl₃): δ = 43.1 (1C, N-CH₃), 45.5 (2C, N-
(CH₃)₂), 52.4 (2C, OCH₃), 56.5 (2C, N-CH₂), 58.9 (2C, NCH₂),
25 62.4 (2C, C_{q,Alkyl}), 73.8 (2C, NCH), 122.9, 123.9, 136.3, 149.2
(8C, Ar-C), 158.6 (2C, Ar-C_q), 168.6 (2C, ester), 207.2 (1C,
C=O).

Preparation of complex 1

30 2 mmol of metal salt (FeCl₂) dissolved in 1 ml methanol is
added to 2 mmol of ligand dissolved in 1 ml acetotrile.

- 32 -

After 24 h stirring at RT the solution is concentrated to 0.5 ml total volume and treated with 5 mL of ethylacetate. The solution is sonicated in an ultrasonic bath. The resulting solid is filtered and dried in high vacuum.

5

$\text{FeCl}(\text{N}_2\text{Py}_2\text{EtNMe}_2)\text{Cl}$ $\text{C}_{26}\text{H}_{35}\text{Cl}_2\text{FeN}_5\text{O}_6 \cdot \text{H}_2\text{O}$ $M = 640.34$
g/mol

Analytical data:

10

CHN Analysis	calc. (%)	C 48.77	H 5.51	N 10.94
	found (%)	C 49.15	H 5.79	N 10.61

FAB⁺MS (NBA): 604.2 $[\text{FeCl}(\text{N}_2\text{Py}_2\text{EtNMe}_2 \cdot \text{H}_2\text{O})]\text{H}^+$.

15

Magnetic moment: $\mu = 5.3$ B.M

Redox potential: $E_{1/2}$: 847 mV in acetonitrile

20 IR [cm^{-1}]: 3136 (m, OH), 3094 (m), 2976 (m), 1716 (s), 1600 (m), 1472 (m), 1426 (m), 1274 (s), 784 (m), 648 (w).

UV-Vis (MeOH): 402 nm ($\epsilon = 1651 \text{ cm}^2 \text{ mol}^{-1}$), 313 nm ($\epsilon = 925 \text{ cm}^2 \text{ mol}^{-1}$), 250 nm ($\epsilon = 5123 \text{ cm}^2 \text{ mol}^{-1}$), 219 nm ($\epsilon = 4354 \text{ cm}^2 \text{ mol}^{-1}$).

25

BLEACHING EXPERIMENTS (air mode)

In an aqueous solution containing 10 mM carbonate buffer (pH 10) with 0.6 g/l NaLAS (linear alkylbenzene sulphonate tomato-soya oil or curry-soya oil stained cloths were added and kept in contact with the solution whilst agitating for

30

- 33 -

30 minutes at 30°C. Comparative experiments were performed using 10 µM of the metal complex referred to in the table below.

5 After the wash, the cloths were rinsed with water and subsequently dried at 30°C and the change in colour was measured immediately after drying with a Linotype-Hell scanner (ex Linotype) (t=0 in the table). The tomato stains
10 were left for 24 h in the dark and measured again (t=1 in the table). The change in colour (including bleaching) is expressed as the ΔE value versus white; a lower ΔE value means a cleaner cloth. The measured colour difference (ΔE) between the washed cloth and the unwashed cloth is defined as follows:

15

$$\Delta E = [(\Delta L)^2 + (\Delta a)^2 + (\Delta b)^2]^{1/2}$$

wherein ΔL is a measure for the difference in darkness between the washed and unwashed test cloth; Δa and Δb are
20 measures for the difference in redness and yellowness respectively between both cloths. With regard to this colour measurement technique, reference is made to Commission International de l'Eclairage (CIE); Recommendation on Uniform Colour Spaces, colour difference equations,
25 psychometric colour terms, supplement no 2 to CIE Publication, no 15, Colorimetry, Bureau Central de la CIE, Paris 1978. The results are shown below in the tables.

- 34 -

Tomato oil (TOL)/pH10 with 0.6 g/l NaLAS

	(t=0)	(t=1)
Blank	20	20
FeMeN4pyCl2	10	5
Complex 1	11	6

Curry oil (COL)/pH10 with 0.6 g/l NaLAS

	(t=0)
Blank	54
FeMeN4pyCl2	46
Complex 1	41

5

The experiments presented in the tables above show that the bispidon ligand carrying a tert-amine moiety provides an advantage.


10

- 35 -

CLAIMS:

1. A bleaching composition comprising:

- 5 a) a monomer ligand, L, or transition metal catalyst thereof of a ligand having the formula (I):



- 10 wherein at least one of R1 and R2 is an optionally substituted tertiary amine of the form -C2-C4-alkyl-NR7R8, in which R7 and R8 are independently selected from the group consisting of straight chain, branched or cyclo C1-C12 alkyl, benzyl, the -C2-C4-alkyl- of the -C2-C4-alkyl-NR7R8
- 15 may be substituted by 1 to 4 C1-C2-alkyl, or may form part of a C3 to C6 alkyl ring, and in which R7 and R8 may together form a saturated ring containing one or more other heteroatoms, the other of R1 and R2 being independently selected from:
- 20 -C2-C4-alkyl-NR7R8 as defined above,
 -C1-C24-optionally substituted-alkyl,
 -C6-C10-aryl, -C1-C4-alkyl-C6-C10-aryl,
 a heterocycloalkyl: selected from the group consisting of:
 pyrrolinyl, pyrrolidinyl, morpholinyl, piperidinyl,
- 25 piperazinyl, hexamethylene imine, 1,4-piperazinyl, tetrahydrothiophenyl, tetrahydrofuranyl, tetrahydropyranyl,

- 36 -

and oxazolidinyl, wherein the heterocycloalkyl may be connected to the ligand via any atom in the ring of the selected heterocycloalkyl,

a -C1-C6-alkyl-heterocycloalkyl, wherein the

5 heterocycloalkyl of the -C1-C6-heterocycloalkyl is selected from the group consisting of: piperidinyl, piperidine, 1,4-piperazine, tetrahydrothiophene, tetrahydrofuran, pyrrolidine, and tetrahydropyran, wherein the

10 heterocycloalkyl may be connected to the -C1-C6-alkyl via any atom in the ring of the selected heterocycloalkyl,

a -C1-C6-alkyl-heteroaryl, wherein the heteroaryl of the -C1-C6-alkylheteroaryl is selected from the group consisting of: pyridinyl, pyrimidinyl, pyrazinyl, triazolyl, pyridazinyl, 1,3,5-triazinyl, quinolinyl, isoquinolinyl,

15 quinoxalinyl, imidazolyl, pyrazolyl, benzimidazolyl, thiazolyl, oxazolidinyl, pyrrolyl, carbazolyl, indolyl, and isoindolyl, wherein the heteroaryl may be connected to the -C1-C6-alkyl via any atom in the ring of the selected heteroaryl and the selected heteroaryl is optionally
20 substituted by -C1-C4-alkyl, -C0-C6-alkyl-phenol, -C0-C6-alkyl-thiophenol, -C2-C4-alkyl-thiol, -C2-C4-alkyl-thioether, -C2-C4-alkyl-alcohol, -C2-C4-alkyl-amine, and a -C2-C4-alkyl-carboxylate;

25 R3 and R4 are independently selected from hydrogen, C1-C4-alkyl, phenyl, electron withdrawing groups and reduced products and derivatives thereof;

X is selected from: C=O, a ketal derivative of C=O, a

30 thioketal of derivative of C=O, and $-[C(R_6)_2]_y-$ wherein y takes a value 0 or 1; each R6 is independently selected


- 37 -

from hydrogen, hydroxyl, O-C1-C24-alkyl, O-benzyl, O-(C=O)-C1-C24-alkyl, C1-C24-alkyl;

z groups are same heteroaromatic groups, selected from the group consisting of: pyridinyl; pyrimidinyl; pyrazinyl; triazolyl; pyridazinyl; 1,3,5-triazinyl; quinolinyl; isoquinolinyl; quinoxalinyl; imidazolyl; pyrazolyl; benzimidazolyl; thiazolyl; oxazolidinyl; pyrrolyl; carbazolyl; indolyl; and isoindolyl, and the selected Z is optionally substituted by -C1-C4-alkyl;

b) the balance carriers and adjunct ingredients.

2. A bleaching composition according to claim 1, wherein z



is , wherein R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkyl-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl.

3. A bleaching composition according to claim 2, wherein R is H or -C1-C4-alkyl.

4. A bleaching composition according to claim 3, wherein R is H.


5. A bleaching composition according to claim 1, wherein z is selected from the group consisting of: benzimidazole, thiazole, and imidazole.

- 38 -


6. A bleaching composition according to any preceding claim, wherein one of R1 and R2 is -CH3.

7. A bleaching composition according to any preceding claim, wherein the -C2-C4-alkyl-NR7R8 is selected from the group consisting of: -CH2CH2-NR7R8, -CH2CMe2-NR7R8, -CMe2CH2-NR7R8, -CMeHCH2-NR7R8, -CMeH-CMeH-NR7R8, -CH2CMeH-NR7R8, -CH2CH2CH2-NR7R8, -CH2CH2CMe2-NR7R8, -CH2CMe2CH2-

NR7R8, -CH2CH2-NEt2, -CH2CH2-N(*i*-Pr)2,



, and



10

8. A bleaching composition according to any preceding claim, wherein X is selected from: C=O, and -[C(R6)2] wherein each R6 is independently selected from hydrogen, hydroxyl, C1-C24-alkoxy and C1-C24-alkyl.

9. A bleaching composition according to any preceding claim, wherein X, is selected from C=O, C(OH)2, *syn*-CH(OH) and *anti*-CH(OH).

20

10. A bleaching composition according to any preceding claim, wherein R7 and R8 are independently selected from the group consisting of -CH3, -C2H5, -C3H7, -C4H9, -C5H11, -C6H13, and -CH2C6H5.

25


- 39 -

11. A bleaching composition according to any preceding claim, wherein at least one of R7 and R8 is an optionally substituted alkyl chain of at least five carbon atoms.


5

12. A bleaching composition according to claim 7, wherein R7 and R8 are $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$ or together form a optionally substituted cyclic structure

selected from the group consisting of:



and







10

13. A bleaching composition according to any preceding claim, wherein R1 is a C2-C4-alkyl-NR7R8.


14. A bleaching composition according to any preceding claim, wherein R1 and R2 are independently C2-C4-alkyl-NR7R8.

15. A bleaching composition according to any preceding claim, wherein -NR7R8 is selected from group consisting

of:




, and




- 40 -

16. A bleaching composition according to any preceding claim, wherein R3 and R4 are selected from the group consisting of: -C(O)O-C1-C24-alkyl, -CH2OC(O)C1-C20-alkyl, benzyl ester, phenyl, benzyl, CN, hydrogen, methyl, and C1-
5 C4-OR wherein R is selected from the group consisting of H, C1-C24-alkyl or C(O)-C1-C24-alkyl.
17. A bleaching composition according to any preceding claim, wherein: R3 = R4.
- 10 18. A bleaching composition according to any preceding claim, wherein R3 and R4 are selected from the group consisting of -CH2OH, and -C(O)O-C1-C6-alkyl.
- 15 19. A bleaching composition according to any preceding claim, wherein R3 and R4 are selected from the group consisting of: -C(O)-O-CH3, -C(O)-O-CH2CH3, and CH2OH.
- 20 20. A bleaching composition according to any preceding claim, wherein Y = 1.
21. A bleaching composition according to any preceding claim, wherein X selected from the group consisting of: C=O, CH2, C(OH)2, syn-CHOR and anti-CHOR, wherein R is H, C1-C24-
25 alkyl or C(O)-C1-C24-alkyl.
22. A bleaching composition according to any preceding claim, wherein X is C=O or C(OH)2.
- 30 23. A bleaching composition according to claim 1, wherein the ligand is:

- 41 -



wherein -NR₆R₇ is selected from the group consisting of -



- 5 24. A bleaching composition according to claims 1 to 23, wherein the complex is of the general formula (A1):



10 in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe(II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI);

15 X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;

20 a represents an integer from 1 to 10;

k represents an integer from 1 to 10;

n represents an integer from 0 to 10;

m represents zero or an integer from 1 to 20; and

- 42 -

L represents a ligand as defined in claims 1 to 22, or its protonated or deprotonated analogue.

25. A bleaching composition according to claim 24, wherein

5 M represents a metal selected from Fe(II)-(III)-(IV)-(V).

26. A bleaching composition according to claim 25, wherein

M represents a metal selected from Fe(II) and Fe(III).

10 27. A bleaching composition according to claim 26, wherein the ligand is present in the form selected from the group consisting of $[\text{FeLCl}]\text{Cl}$ and $[\text{FeL}(\text{H}_2\text{O})](\text{BF}_4)_2$.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/05217

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D3/39 C11D3/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 02 48301 A (UNILEVER PLC ; LEVER HINDUSTAN LTD (IN); UNILEVER NV (NL)) 20 June 2002 (2002-06-20) claims 1-20	1-4, 6-9, 12-27
P, X	WO 02 081613 A (UNILEVER PLC ; LEVER HINDUSTAN LTD (IN); UNILEVER NV (NL)) 17 October 2002 (2002-10-17) page 9, line 11 -page 12, line 8	1-4, 6-9, 12-27
P, X	WO 02 50229 A (UNILEVER PLC ; LEVER HINDUSTAN LTD (IN); UNILEVER NV (NL)) 27 June 2002 (2002-06-27) page 7, line 10 -page 9, line 4 -/-	1-4, 6-9, 12-27

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *8* document member of the same patent family

Date of the actual completion of the international search

11 September 2003

Date of mailing of the international search report

17/09/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3016

Authorized officer

Richards, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/05217

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 072746 A (UNILEVER PLC ; LEVER HINDUSTAN LTD (IN); UNILEVER NV (NL)) 19 September 2002 (2002-09-19) page 15, line 27 -page 18, line 24 -----	1-4, 6-9, 12-27
P,X	WO 02 068574 A (UNILEVER PLC ; LEVER HINDUSTAN LTD (IN); UNILEVER NV (NL)) 6 September 2002 (2002-09-06) page 98, line 11 -page 99, line 32 -----	1-4, 6-9, 12-27
A	WO 00 60045 A (PERKINS CHRISTOPHER MARK ; PROCTER & GAMBLE (US)) 12 October 2000 (2000-10-12) cited in the application claims 1-10 -----	1-27

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 03/05217

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0248301	A	20-06-2002	AU 3318702 A	24-06-2002
			CA 2429629 A1	20-06-2002
			WO 0248301 A1	20-06-2002
			US 2002149000 A1	17-10-2002
WO 02081613	A	17-10-2002	WO 02081613 A1	17-10-2002
			US 2003036492 A1	20-02-2003
WO 0250229	A	27-06-2002	AU 1703002 A	01-07-2002
			CA 2429996 A1	27-06-2002
			WO 0250229 A1	27-06-2002
			GB 2385866 A	03-09-2003
			US 2002169096 A1	14-11-2002
WO 02072746	A	19-09-2002	WO 02072746 A1	19-09-2002
			US 2003054968 A1	20-03-2003
			CA 2431006 A1	19-09-2002
			WO 02072747 A1	19-09-2002
			US 2003054967 A1	20-03-2003
WO 02068574	A	06-09-2002	WO 02068574 A1	06-09-2002
			US 2002198127 A1	26-12-2002
WO 0060045	A	12-10-2000	AU 4061900 A	23-10-2000
			WO 0060045 A1	12-10-2000